

**OKLAHOMA DEPARTMENT OF CORRECTIONS**  
**Initial Physical Examination and Laboratory Requirement: Female**

Initial Physical Examination and Laboratory Requirement: Female (5-6A-4365M (b#3))	
Initial PE Age 20 and younger (5-5B-4312)	<ul style="list-style-type: none"> <li>• Cholesterol, Total, CMP and CBC</li> <li>• Hepatitis C Profile (CPL 162) if known risk factors/exposure, and/or meets certain medical condition per MSRM 140137.06 "Management of Hepatitis C"</li> <li>• HCV Antibody with Reflex RNA (CPL 4677) "Once in lifetime and annually in those with continued high-risk behaviors"</li> <li>• Tuberculin skin test (5-6A-4365M (b#3))</li> <li>• HIV testing – Enzyme-Linked Immunosorbent Assay (ELISA). The Western Blot method will be utilized if the baseline ELISA test result is reactive (or positive). (5-6A-4354M. 5-6A4365M (b#3))</li> <li>• Rapid Plasma Reagin (Syphilis) (CPL # 3503 RPR Reflex to T. Pallidum – PA)</li> <li>• Chlamydial testing (urine) and Gonorrhea testing (urine)</li> <li>• Urine pregnancy testing (if clinically indicated)</li> </ul>
Initial PE Age 21 – 39	<ul style="list-style-type: none"> <li>• Cholesterol, Total, CMP and CBC</li> <li>• Hepatitis C Profile (CPL 162) if known risk factors/exposure, and/or meets certain medical condition per MSRM 140137.06 "Management of Hepatitis C"</li> <li>• HCV Antibody with Reflex RNA (CPL 4677) "Once in lifetime and annually in those with continued high-risk behaviors"</li> <li>• Tuberculin skin test (5-6A-4365M (b#3))</li> <li>• HIV testing – Enzyme-Linked Immunosorbent Assay (ELISA). The Western Blot method will be utilized if the baseline ELISA test result is reactive (or positive). (5-6A-4354M. 5-6A4365M (b#3))</li> <li>• PAP testing</li> <li>• Rapid Plasma Reagin (Syphilis) (CPL # 3503 RPR Reflex to T. Pallidum – PA)</li> <li>• Chlamydial testing and Gonorrhea testing</li> <li>• Urine pregnancy testing (if clinically indicated)</li> </ul>
Initial PE Ages 40-49 yrs.	<ul style="list-style-type: none"> <li>• Lipid Profile, CMP and CBC</li> <li>• Hepatitis Profile (CLP 162) if known risk factors/exposure, meets certain medical conditions or if born 1946-1964 per MSRM 140137.06 "Management of Hepatitis C"</li> <li>• HCV Antibody with Reflex RNA (CPL 4677) "Once in lifetime and annually in those with continued high-risk behaviors"</li> <li>• Mammography</li> <li>• Tuberculin skin test (5-6A-4365M (b#3))</li> <li>• HIV testing – Enzyme-Linked Immunosorbent Assay (ELISA). The Western Blot method will be utilized if the baseline ELISA test result is reactive (or positive). (5-6A-4354M. 5-6A4365M (b#3))</li> <li>• PAP testing</li> <li>• Rapid Plasma Reagin (Syphilis) (CPL # 3503 RPR Reflex to T. Pallidum – PA)</li> <li>• Chlamydial testing and Gonorrhea testing</li> <li>• Urine pregnancy testing (if clinically indicated)</li> </ul>
Initial PE Ages 50-64 yrs.	<ul style="list-style-type: none"> <li>• Lipid Profile, CMP and CBC</li> <li>• Hepatitis Profile (CLP 162) if known risk factors/exposure, meets certain medical conditions or if born 1946-1964 per MSRM 140137.06 "Management of Hepatitis C"</li> <li>• HCV Antibody with Reflex RNA (CPL 4677) "Once in lifetime and annually in those with continued high-risk behaviors "Fecal occult blood testing or at receiving facility"</li> <li>• Mammography</li> <li>• Tuberculin skin test (5-6A-4365M (b#3))</li> <li>• HIV testing – Enzyme-Linked Immunosorbent Assay (ELISA). The Western Blot method will be utilized if the baseline ELISA test result is reactive (or positive). (5-6A-4354M. 5-6A4365M (b#3))</li> <li>• PAP testing</li> <li>• Rapid Plasma Reagin (Syphilis) (CPL # 3503 RPR Reflex to T. Pallidum – PA)</li> <li>• Chlamydial and Gonorrhea testing</li> <li>• Urine pregnancy testing (if clinically indicated)</li> </ul>
Initial PE Ages 65 and older	<ul style="list-style-type: none"> <li>• Lipid Profile, CMP and CBC</li> <li>• Hepatitis C Profile (CPL 162) if known risk factors/exposure, and/or meets certain medical condition per MSRM 140137.06 "Management of Hepatitis C"</li> <li>• HCV Antibody with Reflex RNA (CPL 4677) "Once in lifetime and annually in those with continued high-risk behaviors"</li> <li>• Fecal occult blood testing or at receiving facility</li> <li>• Mammography</li> <li>• Urinalysis</li> <li>• Tuberculin skin test (5-6A-4365M (b#3))</li> <li>• HIV testing – Enzyme-Linked Immunosorbent Assay (ELISA). The Western Blot method will be utilized if the baseline ELISA test result is reactive (or positive). (5-6A-4354M. 5-6A4365M (b#3))</li> <li>• PAP testing</li> <li>• Rapid Plasma Reagin (Syphilis) (CPL # 3503 RPR Reflex to T. Pallidum – PA)</li> <li>• Chlamydial testing and Gonorrhea culture</li> <li>• EKG</li> </ul>