Periodic Physical/ Inmates Returning From GPS Nursing Interventions - Female

MONITORING OF PERIODIC PHYSICALS USES A COLLABORATIVE MULTIDISCIPLINARY TEAM APPROACH. PERIODIC PHYSICAL NURSING INTERVENTIONS ARE APPROVED BY THE CHIEF MEDICAL OFFICER FOR NURSING TO PERFORM IN ACCORDANCE WITH THE PERIODIC PHYSICAL EXAMINATION AND LABORATORY REQUIREMENTS. THESE APPROVED PERIODIC PHYSICAL NURSING INTERVENTIONS DO NOT REQUIRE A HEALTH CARE PROVIDERS ORDER. THE PERIODIC PHYSICAL NURSING INTERVENTIONS ARE BASED UPON NATIONALLY RECOGNIZED EVIDENCE-BASED GUIDELINES AND RECOMMENDATIONS.

Problems:				
Health Care Provider:				
Chronic Clinic labs required: Yes No If "Yes" complete the "Chronic Clinic Nursing Interventions".				
Periodic Physical Laboratory Requirements - Female:				
Ages 17-20	Ages 21 - 39	Ages 40 - 49	Ages 50-64	Ages 65 & older
None	None	☐ Fasting Lipid Profile - Every 1-3 years ☐ CMP – Every 1-3 years ☐ Hepatitis C Profile (CPL 162) – Baseline only	□ Fasting Lipid Profile - Every 1-3 years. □ CMP – Annually □ Hepatitis C Profile (CPL 162) – Baseline only if not previously done □ Fecal occult blood testing X 3 beginning age 50 –Annually until age 75, then by providers discretion.	☐ Fasting Lipid Profile - Annually☐ CMP – Every 3 years☐ Fecal occult blood testing X 3 – Annually until age 75, then by providers discretion.
Inmates Returning From GPS Laboratory Requirements – Female:				
	39 & Younger	Initial PE - Ages 40-49 yrs.	Initial PE - Ages 50-64 yrs.	Initial PE - Ages 65 and older
□ Tuberculin skin test □ HIV testing - Enzyme-Linked Immunosorbent Assay (ELISA). The Western Blot method will be utilized if the baseline ELISA test result is reactive (or positive). □ Rapid Plasma Reagin (Syphilis) □ CMP □ Urine pregnancy test (if clinically indicated)		☐ Tuberculin skin test ☐ HIV testing - Enzyme-Linked Immunosorbent Assay (ELISA). The Western Blot method will be utilized if the baseline ELISA test result is reactive (or positive). ☐ Rapid Plasma Reagin (Syphilis) ☐ CMP ☐ Urine pregnancy test (if clinically indicated)	☐ Tuberculin skin test ☐ HIV testing - Enzyme-Linked Immunosorbent Assay (ELISA). The Western Blot method will be utilized if the baseline ELISA test result is reactive (or positive). ☐ Rapid Plasma Reagin (Syphilis ☐ CMP ☐ Urine pregnancy test (if clinically indicated)	 ☐ Tuberculin skin test ☐ HIV testing - Enzyme-Linked Immunosorbent Assay (ELISA). The Western Blot method will be utilized if the baseline ELISA test result is reactive (or positive). ☐ Rapid Plasma Reagin (Syphilis) ☐ CMP
Interventions	:			
□ Inmate identified and procedure explained. □ Hands washed/sanitized and gloves applied. □ Inmate arm positioned, tourniquet applied, puncture site identified and cleansed. □ Venipuncture site: □ Needle gauge size: □ 20 □ 21 □ 22 □ 23 □ 24 □ 25 □ Butterfly □ Huber □ Number of attempts: □ 1 □ 2 □ Other: □ Unable to obtain. Reason: □ Notified: □ Pressure applied, no bleeding, swelling or redness observed. □ Applied: (check all that apply) □ Band-Aid □ Gauze □ Cotton ball □ Tape □ Coban □ Sharps disposed into biohazardous containe.r □ FOBT issued x 3 with instructions.				
 EKG performed. Results placed in EHR. Co-signed to provider. Education/Intervention: Instructed patient to follow-up sick call if experiencing any signs and symptoms that warrant treatment. Inmate verbalizes understanding of instructions. 				
Progress Note:				
RN/LPN Sign	ature/Credentia	ıls:	Dat	e: Time:
Inmate Name (Last, First)				