

## State of Oklahoma **Oklahoma Health Care Authority Zepatier™ Initiation Prior Authorization Form**

Member Name:	Date of Birth:	Member ID#:
		Pharmacy Fax:
Pharmacy Name:Pharmacist Name:		
Prescriber NPI:	Prescriber Name:	Specialty:
		Drug Name:
NDC:S		
Clinical Information		
HCV Genotype (including subtype):		termined:
2. If the member has genotype 1a, does	es the member have the presence of	termined: of virus with NS5A resistance-associated poly-
morphisms? Yes No		
3. METAVIR Equivalent Fibrosis Stage	e: Testing Type:	
Date Fibrosis Stage Determined:  4. Pre-treatment viral load in the last 1		
4. Pre-treatment viral load in the last 1	2 months: Date Tak	cen:
For METAVIR score of <f1, 2nd="" tes<br="">Prior pre-treatment viral load or anti</f1,>		
Prior pre-treatment viral load or antil 5. Does member have decompensated	t henatic disease or Child-Pugh B of	or C? Yes No
		life expectancy (less than 12 months) that
cannot be remediated by treating He	CV? Yes No	, , ,
		ease specialist, or a transplant specialist with-
in the past 3 months? Yes No		
8. If yes, please include name of speci	alist recommending hepatitis C trea	itment:
9. Has the member been previously tre	ment regimen and reason for failur	 e (relapser, null-responder, partial respond-
er):	ment regimen and reason for failur	c (relapser, null-responder, partial respond-
	Delow (if choosing other, please supply	reference citation to support requested therapy):
☐ Zepatier <sup>™</sup> for 12 weeks	, , , , , , , , , , , , , , , , , , , ,	17
■ Zepatier <sup>™</sup> plus ribavirin for		
Zepatier™ plus ribavirin for	12 weeks	
Other:	- tt	
12. Has the member signed the intent to		
13. Has the member been counseled on the harms of illicit IV drug use and alcohol use and agreed to not use illicit IV drugs or alcohol while on or after they finish hepatitis C treatment? Yes No		
14. Has the member initiated immunization		
15. For women of childbearing potential		
		er) and not planning to become pregnant dur-
	nths of completing treatment	
		monal contraception during treatment and for
member	neung treatment. Please list non-no	rmonal birth control options discussed with
		ughout treatment for ribavirin users
		amazepine, rifampin, St. John's wort, efavi-
		e, nafcillin, ketoconazole, bosentan, etraviri-
ne, elvitegravir/cobicstat/emtricitabir		No
17. Have all other clinically significant issues been addressed prior to starting therapy? Yes No		
18. Will member's ALT levels be monitored prior to initiation, at week 8, and as indicated thereafter? Yes No		
This patient is in need of additional support. I recommend this patient be followed by an OHCA Care Management Nurse. <i>Members must be adherent for continued approval. Treatment gaps of therapy longer than 3 days will result in</i>		
denial of payment for subsequent requests for continued therapy. Refills must be prior authorized.		
Prescriber Signature:		Date:
Has the member been counseled on ap	propriate use of Zepatier™ therapy	
Pharmacist Signature:		Date:
Please do not send in chart notes. Specific informat cessing delays. By signature, the prescriber or p		ary. Failure to complete this form in full will result in pro-
cessing delays. by signature, the prescriber of p	กลากลบระบบกกกร เกษ สมบิงษ์ แบบกกใส่แบบ	กาง สบบนาสเษ.

PLEASE PROVIDE THE INFORMATION REQUESTED AND RETURN TO:

University of Oklahoma College of Pharmacy Pharmacy Management Consultants Product Based Prior Authorization Unit Fax: 1-800-224-4014 Phone: 1-800-522-0114 Option 4

CONFIDENTIALITY NOTICE
This document, including any attachments, contains information which is confidential or privileged. If you are not the intended recipient, be aware that any disclosure, copying, distribution, or use of the contents of this information is prohibited. f you have received this document in error, please notify the sender immediately by telephone to arrange for the return of the transmitted documents or to verify their destruction.