OKLAHOMA DEPARTMENT OF CORRECTIONS
Periodic Physical Examination and Laboratory Requirements
Female

Periodic Physical Examination and Laboratory Requirement: Female (5-ACI-6A-27)		
Age	**Requirements**	**Frequency**
Periodic PE Ages 17-20 yrs.	• Physical Examination	• Every 3 years
Periodic PE Ages 21-39 yrs.	• Physical Examination • PAP testing	• Every 3 years • Every 1-3 years. Provider discretion after three consecutive normal tests or hysterectomy.
Periodic PE Ages 40-49 yrs.	• Physical Examination • Fasting Lipid Profile and CMP • Hepatitis C Profile (CPL 162) • Repeat PAP testing • Mammography	• Every 1-3 years • Every 1-3 years • Baseline Only • Every 1-5 years. Provider discretion after three consecutive normal tests or hysterectomy. • Baseline then every 1-2 years until age 75. Provider discretion for high risk patients.
Periodic PE Ages 50-64 yrs.	• Physical Examination • Fasting Lipid Profile • Hepatitis C Profile (CPL 162) • PAP testing • CMP • Fecal occult blood testing X 3 • Mammography • Baseline EKG-age 50	• Every 1-3 years • Every 1-3 years • Baseline Only if not previously done • Every 1-5 years. Provider discretion after three consecutive normal tests or hysterectomy. • Annually • Annually • Baseline, and then every 1-2 years, or by provider discretion for high risk patients. • Baseline, and then by provider discretion
Periodic PE Ages 65 and older	• Physical Examination • Fasting Lipid Profile • PAP testing • CMP • Fecal occult blood testing X 3 • Mammography • Bone Density Screening	• Annually • Annually • Repeat Pap test is needed only if prior abnormality or other clinical indication. • Annually until 75, and then by provider order (CMP anf Fecal occult blood testing x 3) • Every 1-2 years. Provider discretion after age 75 (Mammography) • Age 65 or equivalent fracture risk. Repeat screening not recommended by USPSTF, may be ordered by provider discretion.

Physical Examination and Laboratory Requirements for Inmates Returning From GPS
Female

GPS Physical Examination and Laboratory Requirement: Female	
Initial PE Age 39 and younger	• Tuberculin skin test • HIV testing – Enzyme-Linked Immunosorbent Assay (ELISA). The Western Blot method will be utilized if the baseline ELISA test result is reactive (or positive). • Rapid Plasma Reagin (Syphilis), Chlamydial testing, Gonorrhea culture • CMP • Urine pregnancy testing (if clinically indicated)
Initial PE Ages 40-49 yrs.	• Tuberculin skin test • HIV testing – Enzyme-Linked Immunosorbent Assay (ELISA). The Western Blot method will be utilized if the baseline ELISA test result is reactive (or positive). • Rapid Plasma Reagin (Syphilis), Chlamydial testing, Gonorrhea culture • CMP • Urine pregnancy testing (if clinically indicated)
Initial PE Ages 50-64 yrs.	• Tuberculin skin test • HIV testing – Enzyme-Linked Immunosorbent Assay (ELISA). The Western Blot method will be utilized if the baseline ELISA test result is reactive (or positive). • Rapid Plasma Reagin (Syphilis), Chlamydial testing, Gonorrhea culture • CMP • Urine pregnancy testing (if clinically indicated)
Initial PE Ages 65 and older	• Tuberculin skin test • HIV testing – Enzyme-Linked Immunosorbent Assay (ELISA). The Western Blot method will be utilized if the baseline ELISA test result is reactive (or positive). • Rapid Plasma Reagin (Syphilis) • Chlamydial testing • Gonorrhea culture • CMP