



Infection Control and Prevention Guidelines for Influenza in Hospitals and Other Acute Care Settings

Modes of Influenza Transmission:

The spread of influenza virus occurs through **droplet transmission**, when respiratory droplets from infected persons **directly** fall within 3-6 feet from the infected person after coughing, sneezing, or talking. These droplets infect the mucus membranes (eyes, nose, or mouth) of others, or contaminate surfaces and items in the infected person's environment. The droplets can be picked up from surfaces and **indirectly** spread to mucus membranes after touching contaminated surfaces.

Droplet transmission differs from **airborne transmission**, which is the ability of tiny respiratory droplets to float in the air and travel further distances. Airborne transmission of influenza viruses is unknown. Therefore airborne precautions should be followed during aerosol-generating procedures such as bronchoscopy, intubation, cardiopulmonary resuscitation (CPR), open airway suctioning, and sputum induction.

Influenza-like illness (ILI) is defined as fever* and a cough and/or sore throat in the absence of another **KNOWN** cause of illness.

*temperature of 100° F (37.8° C) or greater

Control and Prevention of Influenza Viruses in Acute Care Facilities:

1. **Early recognition of patients with influenza-like symptoms** (fever of $\geq 100^{\circ}$ F or 37.8° C and cough and/or sore throat) at first entry into the healthcare system. Prevent spread from the source by the following actions:
 - a. **Place a surgical or procedure mask on any client with influenza-like illness.**
 - If wearing a mask would compromise the client's ability to breathe easily, have them cover their mouth and nose with a tissue when coughing or sneezing.
 - b. **Separate symptomatic persons from others by a least 3-6 feet** as quickly as possible.
 - Place patient into a separate room (such as an exam room) with the door closed.
 - c. **Instruct clients to follow respiratory hygiene, cough etiquette, and hand hygiene.**
 - Cover mouth and nose with a tissue when coughing or sneezing, OR cough or sneeze into the upper sleeve, NOT into one's hands.
 - d. **Provide supplies:** Tissues, waste containers, and hand hygiene materials.
2. Use **standard precautions** when caring for all persons including those with influenza-like illness:
 - a. **Perform hand hygiene** with soap and water or alcohol-based hand product after contact with the patient or potentially contaminated surfaces. **Only use alcohol-based hand rubs when hands are NOT visibly soiled.** Perform hand hygiene:
 - Before having direct contact with patients.
 - After contact with blood, body fluids or excretions, mucus membranes, non-intact skin, and wound dressings. After contact with a patient's intact skin such as checking vital signs or positioning a patient.
 - If hands will be moving from a contaminated body site to a clean body site during patient care.
 - After contact with inanimate objects in the immediate vicinity of the patient.
 - After removing personal protective equipment such as gloves, gowns, or masks.
 - b. **Follow respiratory hygiene and cough etiquette** (see 1.c. above).
 - c. **Use gloves** if hand contact with secretions or contaminated surfaces is anticipated.
 - Change gloves between patients. Failure to do so in an infection control hazard!
 - Put on clean gloves just before touching a patient's mucous membranes, non-intact skin, or contaminated items.
 - Gloves do not replace the need for hand hygiene because gloves may be torn during use, and hands can become contaminated during removal of gloves.
 - Remove gloves and perform hand hygiene when exiting the room.

Control and Prevention of Influenza Viruses in Acute Care Facilities (continued):

3. Use **droplet isolation precautions** in addition to standard precautions when caring for patients with suspected or confirmed influenza until at least 24 hours after their fever is gone (without the use of fever-reducing medicine).
 - a. Place patients into **private rooms** OR cohort patients with like respiratory illnesses together.
 - The patient should wear a surgical mask during transport out of room.
 - Limit healthcare personnel contact to those performing direct patient care.
 - Isolation precautions should be continued until at least 24 hours after their fever is gone (without the use of fever-reducing medicine).
 - b. Use a **surgical or procedure mask** for all routine patient care when entering the patient's room.
 - Remove the mask and perform hand hygiene when exiting the room.
 - c. Wear a **gown** when soiling of clothes or skin with blood, body fluids, secretions, or excretions is anticipated.
 - Change gowns between patients.
 - Remove gowns and perform hand hygiene when exiting the room.

4. Use **airborne precautions during aerosol-generating procedures** (see adjacent box).
 - a. Wear an **N95 respirator (mask)**. These provide a seal around the nose and mouth, and filter out small organisms.
 - Only use the brand and size of N95 mask that was used for the fit-test.
 - Check the fit to insure effectiveness with each use.
 - Remove N95 masks and perform hand hygiene when exiting the room.
 - b. Wear **face protection** such as a face shield, a mask with attached shield, or a mask with goggles.
 - c. Use an **airborne infection isolation room (AIIR)** with negative pressure performing 6 to 12 air exchanges per hour for aerosol-producing procedures if possible.

Aerosol-Generating Procedures:

- Such as bronchoscopies, intubation, CPR, open airway suctioning, and sputum induction.
- When available, perform procedures in an Airborne Infection Isolation Room (AIIR) with negative pressure air handling with 6-12 air changes per hour.
- Healthcare personnel need to wear N95 masks with eye protection (goggles or face shield) during these procedures.

Safe Removal of Personal Protective Equipment (PPE): Follow This Sequence and Do Not Touch the Outside of PPE Items:

1. **Gloves:**
 - a. Grasp outside of glove with opposite gloved hand and peel off.
 - b. Hold removed glove in gloved hand.
 - c. Slide fingers of ungloved hand under remaining glove at wrist.
 - d. Peel second glove off over first glove and discard into trash receptacle.
2. **Goggles or Face Shield:**
 - a. Grasp head band or ear pieces.
 - b. Place in designated receptacle for reprocessing or in waste container.
3. **Gown:**
 - a. Unfasten ties.
 - b. Pull away from neck and shoulders, touching inside of gown only.
 - c. Turn gown inside out.
 - d. Fold or roll into a bundle and discard.
4. **Mask or Respirator:**
 - a. Grasp bottom ties or elastic band, then top and remove without contaminating face.

Environmental Infection Control:

1. Use routine cleaning and disinfection strategies during influenza seasons.
2. Focus on frequently touched surfaces.
3. For further guidance, refer to CDC guidance document on environmental infection control:
<http://www.cdc.gov/mmwr/preview/mmwrhtml/rr5210a1.htm>.

Patient and Visitor Management:

1. Post signage at all building entry points regarding reporting of ILI at the first opportunity so that precautions can be initiated.
1. Post respiratory hygiene/cough etiquette signs and provide supplies to perform these actions.
2. Consider limiting points of entry and/or designate those only for persons with febrile respiratory illnesses.
1. Visitors who have been in contact are a possible source of spread:
 - Limit visitors for patients with suspected or confirmed influenza.
 - Screen visitors before entering hospitals when possible.
 - Limit visitor movement within facility.
 - Offer/require masks, gowns, gloves, and hand hygiene during visits.

Healthcare Personnel Surveillance:

1. Monitor healthcare personnel daily for signs and symptoms of ILI.
2. If an employee becomes ill while at work, they should cease patient care activities, and notify their supervisor and/or the employee health nurse.
3. Healthcare personnel who have had an unprotected exposure to influenza may continue to work as long as they are symptomatic and should contact their primary care provider regarding recommendations for antiviral prophylaxis.

Management of Ill Healthcare Personnel:

1. Instruct all workers to stay home if they are sick with ILI symptoms. Healthcare workers with ILI should be excluded from work for at least 24 hours after they no longer have a fever (defined as $\geq 100^{\circ}$ F or 37.8° C) or signs of a fever, without the use of fever-reducing medicines. Those who work in areas where the patients are considered severely immunocompromised should be considered for temporary reassignment or exclusion from work for 7 days from symptom onset or until symptoms have resolved, whichever is longer.
2. Non-healthcare employees with ILI should stay home and not return to work until at least 24 hours after they are free of fever or signs of a fever without the use of fever-reducing medications.
3. Have time-off and return to work policies and procedures in place for health care providers who are asked to stay home because of fever and respiratory symptoms.