# COMANCHE COUNTY Prescription Opioid Overdose Facts


#### NUMBER OF UNINTENTIONAL OVERDOSE DEATHS BY TYPE OF SUBSTANCE

COMANCHE COUNTY, 2013-2017	
All unintentional overdoses	92
Prescription drugs	55
Prescription opioids	48
Illicit drugs	37
Alcohol	14

#### More than four out of five

unintentional prescription drug overdose deaths in Oklahoma involve at least one prescription opioid.

Overdose deaths involving prescription opioids account for more than all illicit drugs combined.





FROM UNINTENTIONAL PRESCRIPTION OPIOID OVERDOSE

# Most common

substances\* in Comanche County overdose deaths:

Methamphetamine Oxycodone Alcohol Hydrocodone Morphine Alprazolam Fentanyl Cocaine 1,1-Difluoroethane Zolpidem

# **State Facts:**

- From 2013-2017, more than 1,900
  Oklahomans died of an unintentional prescription opioid overdose.
- In 2017, enough opioids were dispensed for every adult in Oklahoma to have the equivalent of 156 hydrocodone 10mg tablets.

\*Opioids in bold




## **Opioid Overdose Fast Facts\*** Comanche County Data (2013-2017)

- There were 48 unintentional prescription opioid overdose deaths in Comanche County.
- Comanche County had the 52th highest death rate in the state.
- $\bullet\,$  Females were 27% more likely to die than males.
- Adults aged 35-54 had the highest rate of death.
- Nearly half (44%) of people who died had a history of mental health problems.
- More than two out of three (69%) people who died had a history of substance abuse.
- Learn the warning signs of an overdose 58% of deaths occurred at a home or apartment.

\*Some rates are based on small numbers and may be unstable.



# **COMANCHE COUNTY**




\*2017 Oklahoma Prescription Monitoring Program data

### Comanche County Prescribing Data\*

- The opioid prescription rate for Comanche County adults was 18% lower than the state rate.
- Adults aged 55-64 had the highest opioid prescription rate.
- The most commonly prescribed opioids were:
  - Hydrocodone
  - Oxycodone
  - Tramadol
  - Codeine
  - Morphine
- 16% of Comanche County adults with an opioid prescription had an overlapping prescription for a benzodiazepine. This combination increases the risk of fatal overdose.

#### Unintentional Prescription Opioid Overdose Death Rates by County of Residence<sup>1</sup>, Oklahoma, 2013-2017



Service Oklahoma State Department of Healt ublication was supported by the Cooperative Agreement Number 6 NU17CE002745 03 funded by the Centers for se Control and Prevention (CDC). Its contents are solely the responsibility of the authors and do not necessarily sent the official views of the CDC or the Department of Health and Human Services. This publication was issued Oklahoma State Department of Health (OSDH), an equal opportunity employer and provider. A digital file has deposited with the Publications Clearinghouse of the Oklahoma Department of Libraries in compliance with n 3.114 of Title 65 of the Oklahoma Statutes and is available for download at www.documents.ok.gov. |

